

AD-A130 438

RAMAN SPECTROSCOPIC STUDY OF MOLECULAR ORIENTATION IN
AGTCNO THIN FILMS(U) BROWN UNIV PROVIDENCE RI DEPT OF
CHEMISTRY E KAMITSOS ET AL. 18 FEB 83 TR-83-01


1/1


UNCLASSIFIED

N00014-75-C-0883

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 13 0438

DTIC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract ORN-N00014-75-C-0883 NR-051-514

TECHNICAL REPORT NO. TR83-01

Raman Spectroscopic Study of Molecular Orientation in AgTCNQ Thin Films

by

E. I. Kamitsos and W. M. Risen, Jr.

Prepared for Publication

in

Journal of Chemical Physics

February 18, 1983

Department of Chemistry

Brown University

Providence, R.I. 02912

DTIC
ELECTE
JUL 18 1983
S A D

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited

83 07 18 038

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TR-83-01	2. GOVT ACCESSION NO. 4D-A130438	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Raman Spectroscopic Study of Molecular Orientation in AgTCNQ Thin Films		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Efstratios Kamitsos and William M. Risen, Jr.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Brown University Providence, Rhode Island 02912		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0883 NR-051-539
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research United States Navy		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS ONR-N00014-75-C-0883 NR-051-539
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE February 18, 1983
		13. NUMBER OF PAGES 24
		14. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Distribution Unlimited; Approved for Public Release		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) charge transfer, thin films, molecular orientation, Raman, TCNQ		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Preferential orientation of TCNQ units in thin AgTCNQ films, which were prepared by thermal treatment of vapor-deposited materials on various substrates, is demonstrated by laser Raman spectroscopy. The polarization dependence of Raman intensities in four scattering arrangements have been obtained by varying the polarization direction of the incident light. The experimental results for Raman intensities as a function of substrate orientation are compared with those predicted for films with TCNQ ⁻ units with various postulated types of orientation relative to the substrate and to each other, and also compared with those		

20. Continued

obtained from AgTCNQ solution. The data show that AgTCNQ thin films are characterized by preferred orientation of the TCNQ⁻ units relative to the substrate at an angle $\approx 45^\circ$. The lack of other orientation in the sampled region is taken to indicate that there are many relatively small domains, within each of which $\approx 45^\circ$, which are not oriented with respect to each other.

Raman Spectroscopic Study of Molecular Orientation in AgTCNQ Thin Films

by

Efstratios I. Kamitsos and William M. Risen, Jr.

Chemistry Department, Brown University

Providence, RI 02912

Received:



SEARCHED	INDEXED
SERIALIZED	FILED
APR 1981	
FBI - BOSTON	
A	

Abstract

Preferential orientation of TCNQ⁻ units in thin AgTCNQ films, which were prepared by thermal treatment of vapor-deposited materials on various substrates, is demonstrated by laser Raman spectroscopy. The polarization dependence of Raman intensities in four scattering arrangements have been obtained by varying the polarization direction of the incident light. The experimental results for Raman intensities as a function of substrate orientation are compared with those predicted for films with TCNQ⁻ units with various postulated types of orientation relative to the substrate and to each other, and also compared with those obtained from AgTCNQ solution. The data show that AgTCNQ thin films are characterized by preferred orientation of the TCNQ⁻ units relative to the substrate at an angle $\theta \approx 45^\circ$. The lack of other orientation in the sampled region is taken to indicate that there are many relatively small domains, within each of which $\theta \approx 45^\circ$, which are not oriented with respect to each other.

Raman Spectroscopic Study of Molecular Orientation in AgTCNQ Thin Films

Efstratios I. Kamitsos and William M. Risen, Jr.

Chemistry Department, Brown University

Providence, RI 02912

Introduction

The synthesis, structure and physical properties of TCNQ charge-transfer salts have been the subject of extensive investigations over the past two decades. It is known from X-ray diffraction studies that the common structural characteristic of these materials is face to face stacking of the TCNQ moieties to form pseudo-one dimensional columns, which may be along one crystallographic axis (1) or tilted relative to a crystallographic axis (2). This directionality of packing of the TCNQ units results in anisotropic extensive physical properties, especially their electrical conductivity and magnetic and optical properties. Such details of the mode of packing as the formation of segregated (3) as opposed to mixed stacks (4) of donor and TCNQ molecules, and the degree of charge transfer from donor to TCNQ (5), determine the absolute values of the extensive properties in different directions without affecting the anisotropic character of the solid state TCNQ salts.

The reported X-ray structural results on TCNQ-compounds were obtained on single crystals of salts prepared by solution methods. We have shown (6,7) that in addition to being preparable by these solution methods of Melby *et al.* (8), CuTCNQ and AgTCNQ can be prepared as thin films on a variety of substrates by a thermal method. In this method, alternating layers of the metal and TCNQ, of the appropriate thicknesses, are vapor-deposited on a substrate and then heated at ca 100°C to cause a direct reaction of metal with TCNQ. The MTCNQ-films obtained that way have the same

spectral features as those prepared by the solution methods, and we have found that they exhibit interesting resonance Raman (9) and photo-transformation phenomena (7).

The crystal structures of CuTCNQ and AgTCNQ are not known, but it is likely that crystals of these salts are characterized by stacks of TCNQ⁻ units, as are the other TCNQ-compounds prepared by similar solution procedures. Since the thin films can have important properties due to their form, it is of interest to investigate whether or not such stacking occurs in them as well or whether the TCNQ⁻ units in thermally prepared thin films are disordered in some way. If preferred orientation of TCNQ⁻ units occurs in thin films, it is expected that they would exhibit interesting directional semiconducting properties. If there is no such order, a new basis for understanding their electronic and optical properties will need to be developed.

Vibrational spectroscopy can be employed to measure the existence and the extent of molecular orientation in a variety of materials. For example Raman depolarization and infrared dichroism measurements have been used to determine the degree of molecular orientation in polymeric fibers (10-12), and to study chain conformations in proteins (13). Recently, we have used polarized Raman scattering to probe the extent of molecular orientation in glass systems. Thus, we showed that $(\text{PO}_3)_n^{-n}$ metaphosphate chains are preferentially oriented with the axis of $(\text{NaPO}_3)_x$ glass fibers drawn from the melt (14), and that boroxol rings are preferentially oriented in response to an applied stress in blown films of B_2O_3 glass (15).

In this study polarized Raman spectroscopy has been employed to explore the existence of preferred orientation of the TCNQ⁻ units in AgTCNQ thin films prepared by the thermal method. The experimental results for Raman intensities as a function of sample orientation are compared with those predicted for films with TCNQ⁻ units with various postulated types of orientation relative to the substrate and to each other.

Experimental

The polarized Raman scattering measurements were performed on AgTCNQ thin films since they are of better optical quality than CuTCNQ thin films and exhibit analogous phenomena. The preparation of thin films has been described previously (6,7). Briefly, films of ca 2000 Å thickness were prepared on three substrates that have different electrical conductivities. A metal, aluminum film (ca 5 μm thick vapor deposited on a fused silica plate), an insulator (a flat fused silica plate), and a semiconductor, Si(100) wafer, were the substrates. To provide a basis for comparison, polarized Raman spectra also were measured on a CH₃CN solution of AgTCNQ, in which the TCNQ⁻ ions are randomly oriented. The AgTCNQ powder from which the dilute solution was made was prepared by the metathetical reaction of AgNO₃ and LiTCNQ (8).

Raman spectra were obtained on a Jarrell-Ash 25-300 laser Raman spectrometer using the 632.8 nm line of a Spectra Physics 125A He-Ne laser. A 90° scattering geometry was used, with Z defined as the direction of propagation of the incident light and X as that of the scattered light. The plane of polarization of the incident light was controlled by a half-wave plate, coated for optimal performance at 632.8 nm. The polarization of the scattered light was determined using a Polaroid sheet analyzer. To compensate for the difference in response of the gratings and other optical components of the spectrometer for different polarizations of the light entering the spectrometer, a scrambler was placed before the entrance slit.

The convention used to describe the polarized Raman spectra, $i(kl)j$, is that of Porto et al. (16). The symbols outside the parentheses to the left and the right are the propagation directions of the incident and scattered light respectively. The symbols inside the parentheses, left to right, are the polarization directions of the incident and scattered light. The Raman depolarization data were obtained for the totally symmetric (Ag) mode of AgTCNQ at ca 1605 cm⁻¹, which is the strongest band of the spectrum for this particular excitation wavelength. The

optical arrangements and spectrometer response were calibrated by measuring the depolarization ratios of CCl_4 . The depolarization ratios obtained are $\rho = 0.010 \pm 0.001$ for the 459 cm^{-1} band, and $\rho = 0.75 \pm 0.01$ for the 314 and 218 cm^{-1} band, in excellent agreement with the established values (17).

Theory

For randomly oriented molecules the power of the Raman scattered radiation is determined by the space-averaged value of the square of the induced dipole moment. Thus, the average of the squares of the derived polarizability tensor components, over all orientations of the molecular-fixed axes (x,y,z) relative to the laboratory-fixed axes (X,Y,Z) are required. Then, the Raman intensities are functions of two quantities, which are invariant under any transformation of coordinate system. These quantities are the mean polarizability $\bar{\alpha}$ and the anisotropy γ , functions of the derived molecular polarizability tensor elements, α_{pq} , where

$$\alpha_{pq} = \left(\frac{\partial^2 \alpha}{\partial p \partial q} \right)_0$$

and

$$\bar{\alpha} = \frac{1}{3}(\alpha_{xx} + \alpha_{yy} + \alpha_{zz}) \quad (1)$$

$$\gamma^2 = \frac{1}{2}[(\alpha_{xx} - \alpha_{yy})^2 + (\alpha_{yy} - \alpha_{zz})^2 + (\alpha_{zz} - \alpha_{xx})^2 + 6(\alpha_{xy}^2 + \alpha_{yz}^2 + \alpha_{zx}^2)] \quad (2)$$

The space averages of the squares of the scattering tensor components, in laboratory coordinates, are calculated to be (18,19):

$$\langle \alpha_{XX}^2 \rangle = \langle \alpha_{YY}^2 \rangle = \langle \alpha_{ZZ}^2 \rangle = \frac{45(\bar{\alpha})^2 + 4\gamma^2}{45} \quad (3)$$

$$\langle \alpha_{XY}^2 \rangle = \langle \alpha_{YZ}^2 \rangle = \langle \alpha_{XZ}^2 \rangle = \frac{\gamma^2}{15} \quad (4)$$

In the case of partial or complete molecular orientation eqn (3) and (4) do not hold and the Raman band intensities depend explicitly on the experimental geometry, the position of the molecular coordinates relative to the laboratory coordinates.

For AgTCNQ thin films we postulate that if there is a preferred orientation of the TCNQ⁻ units relative to the substrate, it is as shown schematically in Figure 1(a). Specifically, this postulate is that each TCNQ⁻ unit is oriented in such a way that the two CN groups at the same end of the molecule are pointing towards the substrate and that the molecular plane makes an angle θ with the plane of the substrate.

In Figure 1(a), x is defined as the long molecular axis, while the short molecular axis y is parallel to the substrate, under the assumptions made here. In Figure 1(b) the molecular axes (x, y, z) are shown relative to the substrate axes (A, B, C), with C defined as being perpendicular to the substrate plane. To specify any coordinate system relative to another three Euler angles are required, and in this paper the Euler angle definitions of Wilson-Decius and Cross are adopted (18). Two of the Euler angles are designated as θ and ϕ in Figure 1(b). The third angle, ψ , is defined as the angle that the y axis makes with the intersection of the (AB) and (xy) planes. Since y is here parallel to the substrate, ψ is zero in Figure 1(b). So the two angles, θ (fixed by the postulate) and ϕ , define completely the (x, y, z) coordinate system relative to the substrate (A, B, C) coordinates. While this postulate is that all TCNQ⁻ units are oriented in the same way relative to the substrate, there is no obvious reason to believe that all TCNQ⁻ units sampled by the source beam are parallel to each other, since relatively large film areas are sampled. It is more likely that the way the film differs from a crystal is that there are domains such that the TCNQ⁻ units in one are parallel to each other but not parallel to units of other domains even though the orientation of each unit relative to the substrate is the same. This is equivalent to saying that θ is fixed but $0 \leq \phi \leq 2\pi$.

In the Raman scattering experiments we can control only the orientation of the substrate relative to the laboratory coordinates in addition to the control of the

polarization. For these measurements the substrate was oriented with its C axis in the (XZ) plane, Figure 1(c). Measurements in two other orientations, where C was in the (XY) or (YZ) planes, also were tried but the measured intensities were too weak to give reliable intensity ratios. So, we consider here only the case where C is in the (XZ) plane. The angle β is experimentally determined and takes values from zero to 90° . The equivalent to angle ϕ (of Figure 1(b)) is zero in Figure 1(c). The third angle, ψ , specifies the variation of A and B axes, within the substrate plane, and is arbitrary. To simplify the calculations we set $\psi = 0$ (Figure 1(c)), but even if ψ ($0 \leq \psi \leq 2\pi$) is included and then averaged over all possible values the expressions obtained for the averages of the squares of the tensor elements (in X,Y,Z coordinates) are the same as are obtained with $\psi = 0$.

Since the intensity of Raman scattered light is directly proportional to the space-averaged value of the square of the relevant α_{pq} element, the derived polarizability tensor elements in (X,Y,Z) coordinates, in terms of derived molecular polarizability elements α_{pq} , are needed. Here P is the direction of polarization of the incident light and Q is that of the scattered light. To obtain these expressions needed, the components of the polarizability tensor in substrate coordinates, α_{FG} , are first expressed in terms of α_{pq} by:

$$\alpha_{FG} = T_1 \alpha_{pq} \tilde{T}_1, \quad (5)$$

where T_1 is the transformation matrix relating the (x,y,z) and (A,B,C) axes systems: $(A,B,C) = T_1(x,y,z)$ and \tilde{T}_1 is its transpose. The elements of matrix T_1 are the direction cosines, that is the cosines axis A makes with axes x, y and z, and so on. The form of matrix T_1 is:

$$T_1 = \begin{pmatrix} \cos\theta \cos\phi & -\sin\phi & \sin\theta \cos\phi \\ \cos\theta \sin\phi & \cos\phi & \sin\theta \sin\phi \\ -\sin\theta & 0 & \cos\theta \end{pmatrix} \quad (6)$$

The α_{PQ} elements are now expressed in terms of α_{FG} by:

$$\alpha_{PQ} = T_2 \alpha_{FG} \tilde{T}_2 \quad (7)$$

Here, T_2 is the transformation matrix relating the (X,Y,Z) and (A,B,C) axes systems:

(X,Y,Z) = T_2 (A,B,C) and is given by:

$$T_2 = \begin{pmatrix} \cos\beta & 0 & \sin\beta \\ 0 & 1 & 0 \\ -\sin\beta & 0 & \cos\beta \end{pmatrix} \quad (8)$$

Combining eqn (5) and (7) results in the following expression for α_{PQ} :

$$\alpha_{PQ} = T_2 T_1 \alpha_{pq} \tilde{T}_1 \tilde{T}_2 = (T_2 T_1) \alpha_{pq} (\tilde{T}_2 \tilde{T}_1) \quad (9)$$

In a tensor form α_{PQ} is written as:

$$\alpha_{PQ} = \begin{pmatrix} \alpha_{XX} & \alpha_{XY} & \alpha_{XZ} \\ \alpha_{YX} & \alpha_{YY} & \alpha_{YZ} \\ \alpha_{ZX} & \alpha_{ZY} & \alpha_{ZZ} \end{pmatrix} \quad (10)$$

If we assume that $TCNQ^-$ retains the molecular symmetry (D_{2h}) of the neutral species, then the polarizability tensor in molecular coordinates and Ag vibrational modes is given as follows (20,21):

$$\alpha_{pq} = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} \quad (11)$$

For the 90° scattering geometry four Raman configurations are possible:

Z(YY)X, Z(YZ)X, Z(XZ)X and Z(XY)X, so the tensor elements α_{YY} , α_{YZ} , α_{XZ} and α_{XY} are needed to compute Raman scattering intensities. By employing eqn (9) and (11) and the expressions for T_1 and T_2 these four elements are calculated. When squared and averaged over all possible values of ϕ , the results are:

$$\langle \alpha_{YY}^2 \rangle = \frac{1}{8} [(a \cos^2 \theta + c \sin^2 \theta + b)^2 + (\sqrt{2} a \cos^2 \theta + \sqrt{2} c \sin^2 \theta)^2 + 2b^2] \quad (12)$$

$$\langle \alpha_{YZ}^2 \rangle = \frac{1}{8} (a \cos^2 \theta + c \sin^2 \theta - b)^2 \sin^2 \theta + \frac{1}{2} (a-c)^2 (\cos^2 \theta \sin^2 \theta) \cos^2 \theta \quad (13)$$

$$\begin{aligned} \langle \alpha_{XZ}^2 \rangle = & \frac{1}{8} [4(a-c)^2 (\sin^2 \theta - \cos^2 \theta)^2 + 4(a \sin^2 \theta + c \cos^2 \theta - b)^2 - \\ & - (a \cos^2 \theta + c \sin^2 \theta - b)^2] \cos^2 \theta \sin^2 \theta + \frac{1}{2} (a-c)^2 (\cos^2 \theta \sin^2 \theta) (\sin^2 \theta - \cos^2 \theta)^2 \quad (14) \end{aligned}$$

$$\langle \alpha_{XY}^2 \rangle = \frac{1}{8} (a \cos^2 \theta + c \sin^2 \theta - b)^2 \cos^2 \theta + \frac{1}{2} (a-c)^2 (\cos^2 \theta \sin^2 \theta) \sin^2 \theta \quad (15)$$

Thus the various $\langle \alpha_{PQ}^2 \rangle$ are now expressed in terms of molecular polarizability elements and trigonometric functions of angles θ and β .

Results and Discussion

AgTCNQ Solution Spectra

Representative polarized Raman spectra of AgTCNQ in CH_3CN solution are shown in Figure 2, for the four different combinations of the polarization of the incident and scattered light. The measured intensities are normalized relative to the intensity of the Z(YY)X orientation, so that three intensity ratios are obtained experimentally. The ratios reported here are the average of at least three sets of experiments; $\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle = 0.33 \pm 0.01$, $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle = 0.34 \pm 0.01$ and $\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle = 0.33 \pm 0.01$. These results are in excellent agreement with the theoretically predicted relative intensities for complete randomness, since they are equal, within experimental error, as predicted by eqns. (4). The three measured ratios give an average of 0.33 ± 0.01 for this Ag mode of TCNQ^- in solution.

For this particular vibrational mode, which is an in-plane, totally symmetric mode (22), it is reasonable to set $c = \alpha_{zz} = 0$. Using $\alpha_{xx} = a$ and $\alpha_{yy} = b$, $\alpha_{ij} = 0$ for $i \neq j$ and combining eqn. (1), (2), (3) and (4) the calculated values of R are:

$$R = \frac{\langle \alpha_{YZ}^2 \rangle}{\langle \alpha_{YY}^2 \rangle} = \frac{\langle \alpha_{XZ}^2 \rangle}{\langle \alpha_{YY}^2 \rangle} = \frac{\langle \alpha_{XY}^2 \rangle}{\langle \alpha_{YY}^2 \rangle} = \frac{a^2 + b^2 - ab}{3a^2 + 3b^2 + 2ab} \quad (16)$$

The ratio R approaches the value $R = \frac{1}{3}$ only when $\frac{a}{b}$ (or $\frac{b}{a}$) is very large and becomes equal to $\frac{1}{3}$ when $a \neq 0$, $b=0$ or $b \neq 0$, $a=0$.

Thus, the solution results suggest that one of the non-zero diagonal elements of the α_{pq} tensor is very large relative to the others. The 632.8-nm line used for Raman excitation is within the red band system of TCNQ^- (23,24), so the 1605 cm^{-1} band is resonantly enhanced. The red band system involves the ${}^2B_{2g} \rightarrow {}^2B_{1u}^{(1)}$ electronic transition, which is x-polarized (23), so it is reasonable to assume that, due to resonance enhancement, the $\alpha_{xx} = a$ element will be much larger than $\alpha_{yy} = b$. Thus, the Raman depolarization data obtained from the $\text{AgTCNQ-CH}_3\text{CN}$ solution spectra show the kind of relative intensities expected for randomly oriented TCNQ^- units, with $a \gg b$.

AgTCNQ Thin Film Spectra.

Representative polarized Raman spectra of AgTCNQ thin films on Al substrate are shown in Figure 3, for different values of angle β . Comparison with the solution spectra of Figure 2, shows that there is a remarkable difference in relative intensities and a strong dependence on angle β . This observation shows that some orientation of the TCNQ^- units relative to the substrate must exist. The experimental relative intensities for different substrates and orientation angle β are given in Table 1.

The comparison of the experimental relative intensities with the ones predicted on the basis of the postulates developed above can now be used to determine the orientation in AgTCNQ thin films. Using the relations derived in the previous section, with $c = 0$ and $a \gg b$, the general expressions given in eqns (12)-(15) can now be simplified to yield:

$$\langle \alpha_{YY}^2 \rangle = \frac{3}{8} a^2 \cos^4 \theta \quad (17)$$

$$\langle \alpha_{YZ}^2 \rangle = \frac{1}{8} a^2 \cos^4 \theta \sin^2 \beta + \frac{1}{2} a^2 (\cos^2 \theta \sin^2 \theta) \cos^2 \beta \quad (18)$$

$$\begin{aligned} \langle \alpha_{XZ}^2 \rangle = & \frac{1}{8} (8 \sin^4 \theta + 3 \cos^4 \theta - 8 \sin^2 \theta \cos^2 \theta) a^2 (\cos^2 \beta \sin^2 \beta) + \\ & + \frac{1}{2} a^2 (\cos^2 \theta \sin^2 \theta) (\sin^2 \beta - \cos^2 \beta)^2 \end{aligned} \quad (19)$$

$$\langle \alpha_{XY}^2 \rangle = \frac{1}{8} a^2 \cos^4 \theta \cos^2 \beta + \frac{1}{2} a^2 (\cos^2 \theta \sin^2 \theta) \sin^2 \beta \quad (20)$$

Since only $\langle \alpha_{YY}^2 \rangle$ is independent of β , the relative intensities with respect to that orientation are obtained.

$$\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle = \frac{1}{3} \sin^2 \beta + \frac{4}{3} \tan^2 \theta \cos^2 \beta \quad (21)$$

$$\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle = \frac{1}{3} (8 \tan^4 \theta - 8 \tan^2 \theta + 3) \cos^2 \beta \sin^2 \beta + \frac{4}{3} \tan^2 \theta (\sin^2 \beta - \cos^2 \beta)^2 \quad (22)$$

$$\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle = \frac{1}{3} \cos^2 \beta + \frac{4}{3} \tan^2 \theta \sin^2 \beta \quad (23)$$

The derived expressions for the theoretical relative intensities are functions of β and θ only. Since θ is fixed, but unspecified by the model, and β is the experimental angle, these relative intensity functions (eqns (21)-(23)) can be plotted versus β , for different values of θ . The calculated curves are shown in Figures (4), (5) and (6), as solid lines. By calculating the first and second derivative (with respect to β) on the right hand side of eqns (21)-(23) it is easy to show that the $\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ and $\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ functions have an inflection point at $\beta = 45^\circ$, for $0 \leq \beta < 90^\circ$. Also the function $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ has inflection points at $\beta = 22.5^\circ$ and $\beta = 67.5^\circ$, while the specific value of angle θ determines whether it shows a maximum or minimum value. Thus for $\theta < 19.88^\circ$ or $\theta > 59.44^\circ$ it shows a maximum at $\beta = 45^\circ$, while for $19.88^\circ < \theta < 59.44^\circ$ the function shows a minimum at $\beta = 45^\circ$.

Finally for $\theta = 19.88^\circ$ or $\theta = 59.44^\circ$ the $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ function does not depend on β and it is a constant determined by θ .

In Figures (4), (5) and (6) we also have plotted the experimental relative intensities for different orientation angle β and various substrates. It is shown that for all three relative intensities there is good agreement between experimental values and calculated curves, especially in the observed trends. The experimental values of $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ relative intensity show well defined minima at $\beta = 45^\circ$, for all three substrates, in excellent agreement with the predicted curves. These experimental values indicate that an angle θ in the 45° to 52.5° range is expected. The other two experimental relative intensities, $\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ and $\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle$, show also the predicted trends, and an orientation angle θ in the 40° to 45° range. Thus, the main experimental result of this investigation is that the predicted relative intensity patterns are observed in the $\theta = 40^\circ$ to 52.5° range.

In Raman depolarization studies of this kind there are several experimental errors and optical effects that can affect, to some extent, the experimental observations. For example, for ratios involving weak bands, noise is one source of error. Low scanning speed and high time constant were used to reduce this source of potential error. Microdefects of the thin films can also cause the incident light to lose its direction of propagation and its initial polarization. Experimental errors, and thus deviations from the predicted results, are expected to be greater for $\beta = 0^\circ$ and 90° , due to the fact that the thin films are substrate-supported and so the interference of the substrate with the incident or scattered light will be a maximum for $\beta = 0^\circ$ and 90° respectively.

The optical effects that are important to consider in these measurements, are refraction and polarization-selective reflection. Since AgTCNQ is an anisotropic material, birefringence or double refraction is expected even at normal incidence.

Thus, for each orientation angle the beam will be refracted and split into two beams with polarization orientations that are mutually orthogonal. The optical indicatrix (index ellipsoid) for these films is not known, so it is impossible to estimate the incident beam deviation or its polarization change, or thus their effect on experimental intensity ratios. The reflected or transmitted light is polarization-selective. For a homogeneous and isotropic medium the transmitted light polarized in the plane of incidence (XZ) is of higher intensity than the transmitted light polarized perpendicularly, (YY), to the plane of incidence (25). If we assume that a fraction of the Raman scattering is excited by the transmitted light, then higher intensity light is available for the (XZ) than the (YY) scattering configuration. This may be the reason that the $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ relative intensities are displaced to higher θ values (40° - 52.5°) compared to $\theta = 40^\circ$ - 45° obtained from the other two relative intensities. AgTCNQ thin films are highly absorbing (with 632.8 nm excitation), so it is expected that most of the measured Raman scattering is generated at the surface or within a small thickness of the film. In this case it is expected that the impact of different optical phenomena on our measurements will be smaller than for transparent materials. In any event, regardless the effect of the optical phenomena on absolute Raman intensities, the effect on the relative intensities is expected to be small, so the observed patterns are the true ones.

In conclusion the data show that AgTCNQ thin films are characterized by preferred orientation of the TCNQ⁻ units relative to the substrate, and that agreement of the experimental results with the calculated relative intensities is obtained with $\theta \approx 45^\circ$. The presence of this (θ) ordering and the lack of other orientation in the sampled region is taken to indicate that there are many relatively small domains, within each of which $\theta \approx 45^\circ$, which are not ordered with respect to each other.

Acknowledgments

This work was supported in part by the Office of Naval Research. The authors are grateful for this and the support and use of facilities, especially the Microelectronics, Electron Microscopic, and Optical Facilities, of the Materials Research Laboratory of Brown University. Discussions with Vincent Mattera of Brown University are gratefully acknowledged.

References

1. M. Konno and Y. Saito, Acta Cryst. B31, 2007 (1975).
2. H. Kobayashi, Y. Ohashi, F. Marumo and Y. Saito, Acta Cryst. B26, 459 (1970).
3. C.J. Fritchie, Jr., Acta Cryst. 20, 892 (1966).
4. H. Kobayashi and J. Nakayama, Bull. Chem. Soc. Japan, 54, 2408 (1981).
5. J.B. Torrance, Acc. Chem. Res. 12, 79 (1979).
6. E.I. Kamitsos, C.H. Tzinis and W.M. Risen, Jr., Solid State Commun. 42, 561 (1982).
7. E.I. Kamitsos and W.M. Risen, Jr., Solid State Commun. 45, 165 (1983).
8. L. R. Melby et al., J. Am. Chem. Soc. 84, 3374 (1962).
9. E.I. Kamitsos and W.M. Risen, Jr., to be published.
10. R.S. Stein, J. Polym. Sci. 31, 335 (1958).
11. D.I. Bower, J. Polym. Sci. B10, 2135 (1972).
12. S.W. Cornell and J.L. Koenig, J. Appl. Phys. 39, 4883 (1968).
13. E.G. Bendit, Biopolymers, 4, 561 (1966).
14. P.J. Miller, G.J. Exarhos and W.M. Risen, Jr., J. Chem. Phys. 59, 2796 (1973).
15. C. Windisch and W.M. Risen, Jr., J. Non-Cryst. Solids, 48, 325 (1982).
16. T.C. Damen, B. Tell and S.P.S. Porto, Phys. Rev. 142, 570 (1966).
17. K. Nakamoto, "Infrared and Raman Spectra of Inorganic and Coordination Compounds", 3rd ed. J. Wiley, New York 1978.
18. E.B. Wilson, J.C. Decius and P.C. Cross, "Molecular Vibrations. The Theory of Infrared and Raman Vibrational Spectra". Dover Publ. Inc. New York 1980.
19. D.A. Long, "Raman Spectroscopy" McGraw-Hill, New York, 1977.
20. R. London, Adv. Phys. 8, 423 (1964).
21. W.M. McClain, J. Chem. Phys. 55, 2789 (1971).
22. R. Bozio, A. Girlando and C. Pecile, J. Chem. Soc. Farad. Trans. II, 71, 1237 (1975).

23. D.A. Lowitz, J. Chem. Phys. 46, 4698 (1967).
24. A. Bieber and J.J. Andre, Chem. Phys. 5, 166 (1974).
25. M. Born and E. Wolf, "Principles of Optics", Fifth ed., Pergamon Press (1975).

Table 1




Experimental Relative Intensities for Different Substrates and Angle θ


θ	$\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$			$\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle$		
	Si	SiO ₂	Al	Si	SiO ₂	Al
90	0.46	0.42	0.42	1.86	2.15	2.04
75	0.46	0.45	0.44	0.81	1.09	0.97
60	0.46	0.48	0.51	0.61	0.81	0.71
45	0.56	0.56	0.61	0.47	0.80	0.56
30	0.58	0.83	0.80	0.49	0.53	0.50
15	1.01	0.91	1.00	0.45	0.51	0.49
0	1.62	1.08	1.10	0.48	0.47	0.38


θ	$\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$		
	Si	SiO ₂	Al
90	3.18	3.05	3.69
75	1.40	2.05	1.83
60	1.02	1.45	1.38
45	0.77	1.12	1.01
30	0.88	1.38	1.28
15	1.67	1.66	1.86
0	2.59	2.25	1.79


Figure Captions


- Figure 1. Definition of the molecular (x,y,z), substrate (A,B,C) and laboratory (X,Y,Z) coordinate system.
- Figure 2. Polarized Raman spectra of AgTCNQ-CH₃CN solution.
- Figure 3. Polarized Raman spectra of AgTCNQ thin films, at different angles θ .
- Figure 4. $\langle \alpha_{YZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ relative intensity. Solid lines represent curves calculated from eqn (21), for different θ values. Points represent experimental values for thin films on various substrates: \circ Si(100), Δ SiO₂ and \square Al.
- Figure 5. $\langle \alpha_{XZ}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ relative intensity. Solid lines represent curves calculated from eqn (22), for different θ values. Points represent experimental values for thin films on various substrates: \circ Si(100), Δ SiO₂ and \square Al.
- Figure 6. $\langle \alpha_{XY}^2 \rangle / \langle \alpha_{YY}^2 \rangle$ relative intensity. Solid lines represent curves calculated from eqn (23), for different θ values. Points represent experimental values for thin films on various substrates: \circ Si(100), Δ SiO₂, and \square Al.














TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139		Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
S. Ruby DOE (STOR) 600 E Street Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. Bernard Spielvogel U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	Dr. Denton Elliott Air Force Office of Scientific Research Bolling AFB Washington, D.C. 20332	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. David Aikens Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. A. P. B. Lever Chemistry Department York University Downsview, Ontario M3J1P3 Canada	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England		Dr. Stanislaw Szpak Naval Ocean Systems Center Code 6343 San Diego, California 95152	1
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362		Dr. Gregory Farrington Department of Materials Science and Engineering University of Pennsylvania Philadelphia, Pennsylvania 19104	
Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, D.C. 20376		Dr. Bruce Dunn Department of Engineering & Applied Science University of California Los Angeles, California 90024	

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
M. L. Robertson Manager, Electrochemical and Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1	Dr. D. Cipris Allied Corporation P. O. Box 3000R Morristown, New Jersey 07960	1
Dr. Micha Tomkiewicz Department of Physics Brooklyn College Brooklyn, New York 11210	1	Dr. M. Philpot IBM Corporation 5600 Cottle Road San Jose, California 95193	1
Dr. Lesser Blum Department of Physics University of Puerto Rico Rio Piedras, Puerto Rico 00931	1	Dr. Donald Sandstrom Washington State University Department of Physics Pullman, Washington 99164	1
Dr. Joseph Gordon, II IBM Corporation K33/281 5600 Cottle Road San Jose, California 95193	1	Dr. Carl Kannewurf Northwestern University Department of Electrical Engineering and Computer Science Evanston, Illinois 60201	1
Dr. Robert Somoano Jet Propulsion Laboratory California Institute of Technology Pasadena, California 91103	1	Dr. Edward Fletcher University of Minnesota Department of Mechanical Engineering Minneapolis, Minnesota 55455	1
Dr. Johann A. Joebstl USA Mobility Equipment R&D Command DRDME-EC Fort Belvoir, Virginia 22060	1	Dr. John Fontanella U.S. Naval Academy Department of Physics Annapolis, Maryland 21402	1
Dr. Judith H. Ambrus NASA Headquarters M.S. RTS-6 Washington, D.C. 20546	1	Dr. Martha Greenblatt Rutgers University Department of Chemistry New Brunswick, New Jersey 08903	1
Dr. Albert R. Landgrebe U.S. Department of Energy M.S. 66025 Forrestal Building Washington, D.C. 20595	1	Dr. John Wassib Kings Mountain Specialties P. O. Box 1173 Kings Mountain, North Carolina 28086	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>	<u>No. 1017-1</u>
Dr. J. J. Brophy University of Utah Department of Physics Salt Lake City, Utah 84112	1	
Dr. Walter Roth Department of Physics State University of New York Albany, New York 12222	1	
Dr. Thomas Davis National Bureau of Standards Polymer Science and Standards Division Washington, D.C. 20234	1	
Dr. Charles Martin Department of Chemistry Texas A&M University	1	
Dr. Anthony Sammells Institute of Gas Technology 3424 South State Street Chicago, Illinois 60616	1	
Dr. H. Tachikawa Department of Chemistry Jackson State University Jackson, Mississippi 39217	1	
Dr. W. M. Risen Department of Chemistry Brown University Providence, Rhode Island	1	

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 510C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 356A

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. Broadhurst Bulk Properties Section National Bureau of Standards U. S. Department of Commerce Washington, D.C. 20234	2	Dr. K. D. Pae Department of Mechanics and Materials Science Rutgers University New Brunswick, New Jersey 08903	1
Naval Surface Weapons Center Attn: Dr. J. M. Augl, Dr. B. Hartman White Oak Silver Spring, Maryland 20910	1	NASA-Lewis Research Center Attn: Dr. T. T. Serofini, MS-49-1 2100 Brookpark Road Cleveland, Ohio 44135	1
Dr. G. Goodman Globe Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1	Dr. Charles H. Sherman Code TD 121 Naval Underwater Systems Center New London, Connecticut 06320	1
Professor Matsuo Ishida Department of Macromolecular Science Case-Western Reserve University Cleveland, Ohio 44106	1	Dr. William Risen Department of Chemistry Brown University Providence, Rhode Island 02191	1
Dr. David Soong Department of Chemical Engineering University of California Berkeley, California 94720		Mr. Robert W. Jones Advanced Projects Manager Hughes Aircraft Company Mail Station D 132 Culver City, California 90230	1
Dr. Curtis W. Frank Department of Chemical Engineering Stanford University Stanford, California 94035		Dr. C. Giori IIT Research Institute 10 West 35 Street Chicago, Illinois 60616	
Picatinny Arsenal Attn: A. M. Anzalone, Building 3401 SMUPA-FR-M-D Dover, New Jersey 07801	1	Dr. R. S. Roe Department of Materials Science and Metallurgical Engineering University of Cincinnati Cincinnati, Ohio 45221	1
Dr. J. K. Gillham Department of Chemistry Princeton University Princeton, New Jersey 08540	1	Dr. Robert E. Cohen Chemical Engineering Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Dr. E. Baer Department of Macromolecular Science Case Western Reserve University Cleveland, Ohio 44106	1	Dr. T. P. Conlon, Jr., Code 3622 Sandia Laboratories Sandia Corporation Albuquerque, New Mexico	1

TECHNICAL REPORT DISTRIBUTION LIST, 356A

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Martin Kaufman Code 38506 Naval Weapons Center China Lake, California 93555	1	Professor C. S. Paik Sung Department of Materials Sciences and Engineering Room 8-109 Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Professor S. Senturia Department of Electrical Engineering Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor Brian Newman Department of Mechanics and Materials Science Rutgers, The State University Piscataway, New Jersey 08854	1
Dr. T. J. Reinhart, Jr., Chief Composite and Fibrous Materials Branch Nonmetallic Materials Division Department of the Air Force Air Force Materials Laboratory (AFSC) Wright-Patterson AFB, Ohio 45433	1	Dr. John Lundberg School of Textile Engineering Georgia Institute of Technology Atlanta, Georgia 30332	1
Dr. J. Lando Department of Macromolecular Science Case Western Reserve University Cleveland, Ohio 44106	1		
Dr. J. White Chemical and Metallurgical Engineering University of Tennessee Knoxville, Tennessee 37916	1		
Dr. J. A. Manson Materials Research Center Lehigh University Bethlehem, Pennsylvania 18015	1		
Dr. R. F. Helmreich Contract RD&E Dow Chemical Co. Midland, Michigan 48640	1		
Dr. R. S. Porter Department of Polymer Science and Engineering University of Massachusetts Amherst, Massachusetts 01002	1		
Professor Garth Wilkes Department of Chemical Engineering Virginia Polytechnic Institute and State University Blacksburg, Virginia 24061	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SOO 5NH United Kingdom	
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Chemistry & Materials Science Department Lawrence Livermore National Lab. Livermore, California 94550	
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auburn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research and Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Johnson Controls 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles and Space Co., Inc. P. O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attn: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1		
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

END

FILMED

8-83

DTIC